# A proposal for spatially consistent weather forecast downscaling via generative deep learning

# David Landry, Anastase Charantonis and Claire Monteleoni

**INRIA** Paris

# **Motivation**

Predicting the state of a sensor network given a weather forecast arises in multiple areas of earth sciences e.g.

- In weather forecast downscaling, when predicting the weather at a station given an NWP forecast.
- In **hydrology**, when predicting streamflow given an NWP forecast.

This problem is challenging because

- NWP is uncertain and provides us with an ensemble forecast.
- NWP models have local biases due to unresolved fine-scale phenomena.

# **Modeling spatial correlations**

Existing downscaling methods treat stations individually. This loses cross-correlations between stations which **are vital in downstream applications**.

**Generative modeling** could allows us to sample the distribution of the full network of stations, but requires new model architectures.

# Use case: Weather Forecast Downscaling

We consider the **weather forecast downscaling** use case. Given the output of a weather forecasting model, what is the likely state of the network of surface weather stations?


To solve this we propose a **cross-attentive transformer** trained within a **denoising diffusion** framework.

Given a weather forecast, how to model the distribution of many in situ measurements and preserve spatial correlations?


# **Preliminary experiments**

We use the transformer architecture for non-generative downscaling of surface temperature an wind gust forecasts.

# **EUPPBench Dataset**



# Sanity check: marginal in situ downscaling




The transformer is equivalent to SOTA for non-generative postprocessing.

The transformer can plausibly be extended for spatially consistent generative modeling, while the DRN cannot needs architectural modifications.

### What does the transformer learn?

We study the **cross-attention** between gridpoints and the Frankfurt/Main during downscaling. Forecast initialized on 2008-02-27T00.


The transformer attends to spatial structures spanning the full domain.




# Prospective generative model

We propose using the **denoising diffusion framework** to sample spatially consistent states for the network of weather stations conditioned on the NWP model output.

The sensor network state progressively goes from a random gaussian distribution (which we can sample) to a denoised, coherent state (which we cannot sample easily).



# Transformer architecture



### Outlook

The transformer network successfully models spatial structures to perform weather forecast downscaling.

The next step is to integrate it into a diffusion framework.

# Challenges and uncertainties

We have a **high-dimensional conditioning** (the NWP forecast) with a **lower-dimensional** target (the station network state), which is unusual.

Evaluation of multivariate ensemble forecasts is still a methodological challenge [Chen2024].

Demonstrate the benefits of a generative approach in downstream applications (hydrology, power production/consumption forecasting).

# References

Ho, J., A. Jain, and P. Abbeel, 2020: **Denoising diffusion probabilistic models.**Proceedings of the 34th International Conference on Neural Information Processing Systems, NIPS '20.

Vaswani, A., N. Shazeer, N. Parmar, J. Uszkoreit, L. Jones, A. N. Gomez, Ł. Kaiser, and I. Polosukhin, 2017: **Attention is All you Need**. Advances in Neural Information Processing Systems.

Demaeyer, J., and Coauthors, 2023: **The EUPPBench postprocessing benchmark dataset v1.0**. Earth System Science Data Discussions, 1–25, https://doi.org/10.5194/essd-2022-465.

Chen, J., T. Janke, F. Steinke, and S. Lerch, 2024: **Generative machine learning methods for multivariate ensemble post-processing.** Ann. Appl. Stat., 18, https://doi.org/10.1214/23-AOAS1784.

Rasp, S., and S. Lerch, 2018: **Neural Networks for Postprocessing Ensemble Weather Forecasts**. Monthly Weather Review, 146, 3885–3900, https://doi.org/10.1175/MWR-D-18-0187.1.

